

目 次

[原 著]

Age and gender differences in body image over the life span:
Relationships between physical appearance, health and functioning
..... Shimpei Ando and Hisao Osada 1

中学生の学校ストレスに対する自己価値の役割
.....西野 泰代 17

日本語版病気認知質問紙の作成と信頼性・妥当性の検討
——血液透析患者による検証——
.....片山富美代・小玉 正博・長田 久雄 28

終末期医療に携わる臨床看護職者のストレスと
ストレス関連成長（Stress-Related Growth）に関する研究
.....逆井 麻利・松田 英子 40

大学生の生活リズム（朝型－夜型）とその認知的評価がストレス反応に及ぼす影響
.....福井 義一・福井 貴子 52

[資 料]

抑うつ対処の自己効力感の向上を目的とした実践に関する効果研究
——大学の授業を活用して——
.....及川 恵・坂本 真士 60

[事例研究]

受診を中断している境界型（IGT）の人における
療養行動の改善を目指した認知の変容の有効性
.....井澤美樹子・伊坂 裕子 67

[会 報]..... 77

会報報告

認定健康心理士資格取得者

学会会則

役員一覧

Contents

Age and gender differences in body image over the life span:
Relationships between physical appearance, health and functioning
Shimpei Ando and Hisao Osada
(Graduate School of Gerontology, J. F. Oberlin University) 1

Self-worth and psychological stress in junior high school students
Yasuyo Nishino
(Graduate School of Education and Development Nagoya University) 17

Development of the Japanese version of the Illness Perception Questionnaire:
Reliability and validity with hemodialysis data
Fumiyo Katayama*¹, Masahiro Kodama*² and Hisao Osada*³
(*¹ Faculty of Engineering, Toin University of Yokohama,
*² Graduate School of Comprehensive Human Sciences, University of Tsukuba,
*³ Graduate School of Gerontology, J. F. Oberlin University) 28

Job stressors and stress-related growth in hospital nurses involved in terminal care nursing
Mari Sakasai*¹ and Eiko Mathuda*¹
(*¹ Faculty of Sociology, Edogawa University) 40

Cognitive appraisal of daily rhythm influences stress reactions among university students
Yoshikazu Fukui*¹ and Takako Fukui*²
(*¹ Tokai-Gakuin University,
*² Gifu SDF Hospital) 52

Psycho-educational university classes promoted self-efficacy for coping with depression
Megumi Oikawa*¹ and Shinji Sakamoto*²
(*¹ Center for the Promotion of Excellence in Higher Education, Kyoto University,
*² College of Humanities and Sciences, Nihon University) 60

Changing cognitions improved self-care behavior in patients
with Impaired Glucose Tolerance who discontinued consultations
Mikiko Izawa*¹ and Hiroko Isaka*²
(*¹ Department of Nursing, Faculty of Health Sciences, Aomori University of Health and Welfare,
*² Nihon University College of International Relations) 67

Official Information 77

Cover design: Haruo Tanabe